

## HOMWORK 10

Due date: Monday of Week 15

Exercises: 1 (a), (b), (c), 6, 10, 12, p.148-149;  
Exercises: 4, 7, 10, 11, 12, 13. p.155-156.

Let  $K$  be a commutative ring with 1. An element  $a \in K$  is called a **unit** if there exists an element  $b \in K$  such that  $ab = 1$ . Denote by  $K^\times$  the set of all units in  $K$ . For example, if  $K$  itself is a field, then  $K^\times = \{x \in K : x \neq 0\}$ . For the integer ring  $\mathbb{Z}$ , we have  $\mathbb{Z}^\times = \{1, -1\}$ . What is  $F[x]^\times$  for a field  $F$ ?

Consider  $\text{Mat}_{n \times n}(K)$  and the following 3 types “elementary matrices” in  $\text{Mat}_{n \times n}(K)$  which are defined as follows. A type I elementary matrix is obtained by multiplying an element  $c \in K^\times$  to a row of  $I_n$ , which is denoted by  $E_n(R_i \leftarrow cR_i)$ . Here as usual,  $I_n$  is the identity matrix. A type II elementary matrix is obtained by adding  $cR_j$  to  $R_i$  of the identity matrix  $I_n$  for some  $c \in K$ , which is denoted by  $E_n(R_i \leftarrow R_i + cR_j)$ . A type III elementary matrix is obtained by switching two rows of  $I_n$ , which is denoted by  $E_n(R_i \leftrightarrow R_j)$ . One can also define elementary row operations similarly and the only difference from what we learned in Chapter I (in that case  $K$  is a field) is: in the first kind elementary row operation, we require that  $c$  is a unit in  $K$  rather than any nonzero elements (we have seen that if  $F$  is a field, a unit in  $F$  is just a nonzero element in  $F$ . The elementary matrices defined here is actually the same as we defined in Chapter I if we generalize “ $F^\times$ ” to “units”.)

Similarly, we can define elementary column operations. We did not even talk about this when  $F$  is a field. The reason for it is: it is unnecessary using column operations for the purpose we did so far. But it is necessary to use elementary elementary column operations when  $K$  is not a field. For a matrix  $j$ , we denote by  $C_j$  its  $j$ -th column. We consider the following 3 types elementary column operations. Type 1,  $e(C_i \leftarrow cC_i)$ , replace  $C_i$  by  $cC_i$  with  $c \in K^\times$ ;  $e(C_i \leftarrow C_i + cC_j)$ : replace  $C_i$  by  $C_i + cC_j$ ;  $e(C_i \leftrightarrow C_j)$ : swap  $C_i$  and  $C_j$ . Then we can also define elementary matrices using elementary column operations applying to  $I_n$ :  $E_n(C_i \leftarrow cC_i)$  ( $c \in K^\times$ );  $E_n(C_i \leftarrow C_i + cC_j)$ ; and  $E_n(C_i \leftrightarrow C_j)$ . For example  $E_n(C_i \leftarrow C_i + cC_j)$  is the matrix obtained by adding  $cC_j$  of  $I_n$  to  $C_i$ .

**Problem 1.** Show that the elementary matrices defined by elementary column operations can be also defined using elementary row operations.

For example  $E_n(C_i \leftarrow cC_i) = E_n(R_i \leftarrow cR_i)$ . It is also easy to check the rest.

**Problem 2.** Show that each elementary matrix in  $\text{Mat}_{n \times n}(K)$  is invertible. Write explicitly each type elementary matrices in  $\text{Mat}_{2 \times 2}(\mathbb{Z})$ .

**Problem 3.** Given a matrix  $A \in \text{Mat}_{m \times n}(K)$ . Let  $e$  be an elementary row operation and let  $E$  be the corresponding elementary matrix. Show that  $e(A) = EA$ . Similarly, if  $e$  is an elementary column operation, and  $E$  is the corresponding elementary matrix. Show that  $e(A) = AE$ .

Just check this case by case. We proved a similar result for elementary row operations in Chapter 1.

If  $K$  is a field, we showed in Chapter 1 that every matrix can be reduced to Row-Reduced Echelon matrix using elementary row operations. If  $K$  is not a field, usually a matrix cannot be reduced to Row-Reduced Echelon matrix in the sense we defined in Chapter 1 using elementary row/column operations. Try to consider what “simple form” you can get for a matrix  $\text{Mat}_{m \times n}(K)$  if  $K = \mathbb{Z}$  or  $F[x]$  using just elementary row operation and elementary column operators. You can define what “simple” means. The next problem will give you some examples.

**Problem 4.** Consider the matrix

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 6 & 6 \end{bmatrix} \in \text{Mat}_{2 \times 3}(\mathbb{Z}).$$

- (1) Using elementary row and elementary column operations (defined above for  $K = \mathbb{Z}$ , namely, in type I,  $c$  is only allowed in  $\mathbb{Z}^\times = \{1, -1\}$ ) to reduce the matrix to

$$B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \end{bmatrix} \in \text{Mat}_{2 \times 3}(\mathbb{Z}).$$

- (2) Find invertible matrices  $P \in \text{GL}_3(\mathbb{Z}), Q \in \text{GL}_2(\mathbb{Z})$  such that  $B = QAP$ .

- (3) Try to use both elementary row and elementary column operations to reduce  $C = \begin{pmatrix} 2 & -1 \\ 1 & 2 \end{pmatrix} \in \text{Mat}_{2 \times 2}(\mathbb{Z})$  to a simple matrix  $C' \in \text{Mat}_{2 \times 2}(\mathbb{Z})$ . Here it is up to you to decide what kind matrices are “simple”. The answer depends on your own interpretation. Is  $C \in \text{GL}_2(\mathbb{Z})$ ?

In the above Problem 4, we only considered matrices with coefficients in  $\mathbb{Z}$ . You can also try similar questions for matrices with coefficients in  $F[x]$  for a field  $F$ . Try to make your own problems and solve them.

Also think about the following question. If  $K = \mathbb{Z}$  or  $F[x]$ . Given  $A \in \text{GL}_n(K)$ , is it possible to reduce  $A$  to the identity matrix using elementary row and column operations? Note that, if this is true, then it will imply that every matrix  $A \in \text{GL}_n(K)$  is still a product of elementary matrices. The answer is Yes if  $K = \mathbb{Z}$  or  $F[x]$ . We will see how to do this in Section 7.4 if  $K = F[x]$ . When  $K = \mathbb{Z}$ , we will see how to do this in a future course (and see how it will be used to determine the structure of finitely generated abelian groups.) Please try some examples using  $2 \times 2$  matrices if  $K = \mathbb{Z}$ . It is indeed easy. If  $K$  is more general, the question is related to something called  $K$ -theory. (The letter “ $K$ ” in  $K$ -theory is not related to the letter “ $K$ ” that we used to denote our ring). Hopefully you will learn something related to  $K$ -theory in the future.

The above problems are no hard at all. They are indeed linear algebra (but over a ring rather than a field).

Do the above problems after Monday’s class.

**Problem 5.** Let  $V = F^2$  and  $W = F^3$ . Compute  $\dim_F \text{Alt}(V^3; F)$  and  $\dim_F \text{Alt}(W^2; F)$ .

You can do Problem 5 after Monday’s class.

Recall that  $S_n$  denotes the symmetric group on  $n$ -elements. In other words,  $S_n$  consists of bijections  $\{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ . Recall that a matrix  $P$  is called permutation matrix if each matrix and each row of  $P$  has only one nonzero element and that nonzero element is 1. (We can view  $P$  as an element of  $\text{GL}_n(K)$  for any commutative ring  $K$  with identity because both 1 and 0 are defined in any such  $K$ . If it is confusing, you can view  $P$  as an element in  $\text{GL}_n(F)$  for a field  $F$ . If it is helpful, you might take  $F$  to be any field you are familiar with). Denote by  $\text{Perm}_n$  the set of all  $n \times n$  permutation matrices. We define

$$e_i = (0, 0, \dots, 0, 1, 0, \dots, 0)^t = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \in F^n$$

with 1 in the  $i$ -th position. For an element  $\sigma \in S_n$ , we consider the matrix  $P_\sigma$  defined by

$$P_\sigma = [e_{\sigma(1)}, e_{\sigma(2)}, \dots, e_{\sigma(n)}].$$

For example, if  $n = 3$ , and  $\sigma \in S_3$  is the element such that  $\sigma(1) = 2, \sigma(2) = 3, \sigma(3) = 1$ , then

$$P_\sigma = [e_{\sigma(1)}, e_{\sigma(2)}, e_{\sigma(3)}] = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}.$$

**Problem 6.** Consider the map  $\theta : S_n \rightarrow \text{Perm}_n$  defined by  $\theta(\sigma) = P_\sigma$ . Show that

- (1)  $\theta(\sigma\tau) = \theta(\sigma)\theta(\tau)$ ;
- (2)  $\theta$  is a bijection;
- (3)  $\det(P_\sigma) = \text{sgn}(\sigma)$ .

Some of these claims were proved in class. For (2), the following fact is useful. Let  $X, Y$  be two finite sets with the same cardinality, and let  $f : X \rightarrow Y$  be an injective map. Then  $f$  must be bijective.

**Problem 7.** Given  $x_1, \dots, x_n \in F$ , consider the matrix

$$A(x_1, x_2, \dots, x_n) = \begin{bmatrix} 1 & x_1 & x_1^2 & \dots & x_1^{n-1} \\ 1 & x_2 & x_2^2 & \dots & x_2^{n-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_n & x_n^2 & \dots & x_n^{n-1} \end{bmatrix}.$$

Compute  $\det(A(x_1, \dots, x_n))$ .

This is a slight generalization of Ex 2, page 155. If you don't know how to do the above problem for general  $n$ , try the case when  $n = 4$ . Do the above two problems after Friday's class.

You don't have to do the next problem. If you have time, try to think about it for some small  $n$  (like  $n = 3, 4$ ). Otherwise, ignore it. Like the authors said on page 162, our focus is not on explicit calculations of determinants of specific matrices.

**Problem 8.** Consider the following  $n \times n$  matrix with coefficients in  $\mathbb{Z}$ :

$$A(n) = \begin{bmatrix} 1 & 2 & 3 & \dots & n \\ 1^3 & 2^3 & 3^3 & \dots & n^3 \\ 1^5 & 2^5 & 3^5 & \dots & n^5 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1^{2n-1} & 2^{2n-1} & 3^{2n-1} & \dots & n^{2n-1} \end{bmatrix}.$$

Compute  $\det(A(n))$ .