HOMEWORK 5

Due date: Oct 30, Monday of Week 10 Exercise: 6, 7, page 66; Exercise: 5, 11, 12, 13, page 73-74;

In the following, we fix a field F.

Problem 1. Let $A \in Mat_{m \times n}(F)$ be a matrix of rank 1.

- (1) Show that there is a matrix $u \in \operatorname{Mat}_{m \times 1}(F)$ and a matrix $v \in \operatorname{Mat}_{1 \times n}(F)$ such that A = uv. (Here uv means the matrix product of u and v).
- (2) If m = n, show that $A^2 = tr(A)A$.

The next problem is a generalization of the above one.

Problem 2. Let $A \in \operatorname{Mat}_{m \times n}(F)$ be a matrix of rank k. Show that there is a matrix $C \in \operatorname{Mat}_{m \times k}(F)$, $R \in \operatorname{Mat}_{k \times n}$ such that A = CR.

(Hint: use elementary row operations. Here one can take R to be the nonzero rows of the RRE matrix of A. What can you say about C? For an integer p with $p < \operatorname{rank}(A)$, is it possible to find matrices $C_1 \in \operatorname{Mat}_{m \times p}(F)$ and $R_1 \in \operatorname{Mat}_{p \times n}(F)$ such that $A = C_1 R_1$? We will go back to this question later.)

Problem 3. Let F be a field of characteristic 0. We view elements of F^3 as column vectors. Consider the matrix

$$A = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 0 & -2 \\ 0 & -1 & 2 \end{bmatrix}$$

and the linear operator $T: F^3 \to F^3$ given by $T(\alpha) = A\alpha$. Compute ker(T) and ker (T^2) . Moreover, show that ker $(T^k) = \text{ker}(T)$ for any positive integer k. Here $T^2 = T \circ T$, and $T^k = T \circ T \circ \cdots \circ T$ (k-times).

Problem 4. Let V be a vector space and let $T: V \to V$ be a linear map. Suppose that $\ker(T) = \ker(T^2)$. Show that $\ker(T^k) = \ker(T)$ for any integer $k \ge 3$.