

### HOMEWORK 3

Due date: Monday of Week 4  
 Exercises: 1, 2, 3, 6, 8, 10, 11, 12, 13, 15, 16, pages 250-251 of Hoffman-Kunze,

The next problem is not closely related to the materials of this week.

**Problem 1.** Given two matrices  $A, B \in \text{Mat}_{n \times n}(\mathbb{C})$ .

- (1) Show that  $\chi_{AB} = \chi_{BA}$ .
- (2) Suppose that  $\deg(\mu_{AB}) > \deg(\mu_{BA})$ , show that  $\mu_{AB} = x\mu_{BA}$ . Here  $x\mu_{BA}$  denote the product of the polynomial  $x$  with the minimal polynomial  $\mu_{BA}$  of  $BA$ .
- (3) Suppose that  $AB$  is diagonalizable, show that  $(BA)^2$  is diagonalizable.
- (4) Give one example such that  $AB$  is diagonalizable but  $BA$  is not diagonalizable.
- (5) Let  $\lambda \in \mathbb{C}, \lambda \neq 0$  and  $r \in \mathbb{Z}$  be a positive integer. Show that  $\dim \text{Ker}(\lambda I - AB)^r = \dim \text{Ker}(\lambda I - BA)^r$ .

The above problem was borrowed from [this link](#). For the first part, see HW 12 of last year. The second part follows from a direct computation. For (3), discuss  $\mu_{BA}$  using (2). For (5), see the proof of Problem 1, HW 12 of last year.

Given a positive integer  $n$ , a partition  $\lambda$  of  $n$  is a sequence of decreasing positive numbers  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$  with  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$  and  $\sum_{i=1}^k \lambda_i = n$ . We write  $\lambda \vdash n$ . Given a sequence of decreasing positive integers  $\lambda = (\lambda_1, \dots, \lambda_k)$  with  $\lambda_1 \geq \dots \geq \lambda_k$ , we also write  $|\lambda| = \sum_{i=1}^k \lambda_i$ . Thus  $\lambda \vdash |\lambda|$ . For example  $(2, 2) \vdash 4, (2, 1, 1) \vdash 4$ . Given a positive integer  $n$ , let  $\mathcal{P}(n)$  be the set of all partitions of  $n$ . Let  $P(n) = |\mathcal{P}(n)|$ , which is the number of all partitions of  $n$ . For example,

$$\mathcal{P}(4) = \{(4), (3, 1), (2, 2), (2, 1, 1), (1, 1, 1, 1)\}.$$

Thus  $P(4) = 5$ .

- Problem 2.**
- (1) Compute  $\mathcal{P}(n)$  and  $|\mathcal{P}(n)|$  for  $n = 5, 6$ .
  - (2) Show that  $P(n)$  is the coefficient of  $x^n$  in the formal power series

$$\prod_{m=1}^{\infty} \left( \frac{1}{1 - x^m} \right) \\
 = (1 + x + x^2 + x^3 + \dots)(1 + x^2 + x^4 + x^6 + \dots)(1 + x^3 + x^6 + x^9 + \dots) \dots$$

Recall that a matrix  $A \in \text{Mat}_{n \times n}(\mathbb{C})$  is called nilpotent if  $A^k = 0$  for some  $k > 0$ . We denote  $\mathfrak{n}_n(\mathbb{C})$  the subset of nilpotent matrices in  $\text{Mat}_{n \times n}(\mathbb{C})$ . On  $\mathfrak{n}_n(\mathbb{C})$ , we define an equivalence relation  $R$  by similarity, namely,  $R = \{(A, B) \in \mathfrak{n}_n(\mathbb{C}) \times \mathfrak{n}_n(\mathbb{C}) : A \text{ is similar with } B\}$ . We consider the equivalence class  $\mathfrak{n}_n(\mathbb{C})/R$ . An element of  $\mathfrak{n}_n(\mathbb{C})/R$  (which is an equivalence class) is called a conjugacy class of a nilpotent matrix. Recall that a typical element in  $\mathfrak{n}_n(\mathbb{C})/R$  is of the form  $\bar{A} = \{B \in \mathfrak{n}_n(\mathbb{C}) : B \text{ is similar with } A\}$  for some  $A \in \mathfrak{n}_n(\mathbb{C})$ . Given a partition  $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$ , we consider the nilpotent matrix

$$A_\lambda = \begin{bmatrix} A_{\lambda_1} & & \\ & \dots & \\ & & A_{\lambda_k} \end{bmatrix} \in \mathfrak{n}_n(\mathbb{C}),$$

where for a positive integer  $m$ ,  $A_m$  denotes the Jordan block with zero in the diagonal of size  $m$ , namely,

$$A_m = \begin{bmatrix} 0 & & & & \\ 1 & 0 & & & \\ & \ddots & \ddots & & \\ & & \ddots & 0 & \\ & & & 1 & 0 \end{bmatrix} \in \text{Mat}_{m \times m}(\mathbb{C}).$$

For example, for  $\lambda = (3, 2) \vdash 5$ , we have

$$A_\lambda = \begin{bmatrix} 0 & 0 & 0 & & \\ 1 & 0 & 0 & & \\ 0 & 1 & 0 & & \\ & & & 0 & 0 \\ & & & 1 & 0 \end{bmatrix}$$

**Problem 3.** Consider the map  $\mathcal{P}(n) \rightarrow \mathfrak{n}_n(\mathbb{C})/R$  defined by  $\lambda \mapsto A_\lambda$ . Show that this map is a bijection.

Given a matrix  $A \in \text{Mat}_{n \times n}(F)$  (for simplicity, we assume that  $F = \mathbb{C}$ ). Let  $c$  be an eigenvalue of  $A$ , we have defined algebraic multiplicity and geometric multiplicity of  $A$  at the eigenvalue  $c$ . We lack standard notations here. Recall that the geometric multiplicity of  $A$  at  $c$  is defined to be  $\dim \text{Ker}(A - cI)$ . Recall that if  $A$  and  $B$  are similar, then they have the same eigenvalues. Moreover, at each eigenvalue, the algebraic multiplicities (and geometric multiplicities) of  $A$  and  $B$  are the same. Conversely, even if  $A$  and  $B$  have the same algebraic and geometric multiplicities at each eigenvalue, it does not mean that  $A$  and  $B$  are similar. A notation which generalize the geometric multiplicity is to consider  $\dim(A - cI)^r$  for every positive integer  $r$ .

**Problem 4.** Given a partition  $\lambda \vdash n$ .

- (1) Give two nilpotent matrices  $A, B$  such that  $A, B$  have the same geometric multiplicity, but  $A$  is not similar to  $B$ .
- (2) Determine  $\dim \text{Ker}(A_\lambda - 0I)^r = \dim \text{Ker}(A_\lambda)^r$  in terms of  $\lambda$  (and  $r$ ).
- (3) Suppose that  $A, B$  are two nilpotent matrices (so the only eigenvalue of  $A, B$  is 0) such that  $\dim \text{Ker}(A)^r = \dim \text{Ker}(B)^r$  for every  $r$ . Is it true that  $A$  is similar to  $B$ ? In other words, if  $A$  is in Jordan canonical form, can the set  $\{\dim \text{Ker}(A)^r : r \geq 1\}$  uniquely determine the corresponding partition of  $A$ ? If so, prove it. If not, give a counter example.

For (2) and (3), if it is hard, at least try the case when  $n = 5$ .

**Problem 5.** Let  $A \in \text{Mat}_{n \times n}(\mathbb{R})$  such that  $(A^2 + 1)^2(A^2 + 2) = 0$ . Find a relatively simple matrix in the conjugacy class of  $A$ .

Hint: This is an exercise from class. You should know what I mean if you attended Wednesday's class.

We consider the finite field  $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$  where  $p$  is a prime integer. For simplicity, we assume that  $p > 2$ . It is known that there is an element  $\kappa \in \mathbb{F}_p^\times = \mathbb{F}_p - \{0\}$  such that  $x^2 - \kappa = 0$  has no solution in  $\mathbb{F}_p$ . For example, if  $p = 3$ , we can take  $\kappa = 2$ ; if  $p = 5$ , we can take  $\kappa = 2$  or  $3$ . Such  $\kappa$  is not unique in general. But if  $\kappa_1, \kappa_2$  are two such numbers, then  $x^2 - \kappa_1\kappa_2^{-1} = 0$  has a solution in  $\mathbb{F}_p$ . For example, if  $p = 5$ ,  $\kappa_1 = 2, \kappa_2 = 3$ , then  $\kappa_1\kappa_2^{-1} = 4$  and  $x^2 - 4 = 0$  has a solution in  $\mathbb{F}_p$ .

**Problem 6.** Fix an element  $\kappa \in \mathbb{F}_p^\times$  such that  $x^2 - \kappa = 0$  has no solution in  $\mathbb{F}_p$ . Show that any element  $g \in \text{GL}_2(\mathbb{F}_p)$  is similar to one of the following type matrices

- (1)  $\begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}, a \in \mathbb{F}_p^\times;$
- (2)  $\begin{bmatrix} a & 1 \\ 0 & a \end{bmatrix}, a \in \mathbb{F}_p^\times;$

$$(3) \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}, a, b \in \mathbb{F}_p^\times, a \neq b;$$

$$(4) \begin{bmatrix} a & b\kappa \\ b & a \end{bmatrix}, a \in \mathbb{F}_p, b \in \mathbb{F}_p^\times.$$

You don't have to submit solutions of the following problem.

- Problem 7.** (1) *Try to classify conjugacy classes of  $\mathrm{GL}_3(\mathbb{F}_p)$ .*  
(2) *Try to classify conjugacy classes of  $\mathrm{GL}_3(\mathbb{C})$  and  $\mathrm{GL}_3(\mathbb{R})$ .*